

For Control of Wild Oats, Ryegrass, Bromes and Broad-leaved Weeds in Winter Wheat, Winter Rye, Winter Triticale
Key Facts

Product Registration Number:	MAPP 18273
Active Ingredient:	70.8 g/Kg pyroxsulam + 14.2 g/Kg florasulam
Pack Size:	1.06 Kg (4 hectares)
Formulation:	Water Dispersible Granule
Maximum Individual Dose:	265 g/ha
Maximum Number of Applications:	1 per crop
Application Timing:	GS11-32 inclusive
Water Volumes:	100-200 Litres water/ha. Optimum 130-150 Litres water/ha
Spray Quality:	FINE to MEDIUM as defined by BCPC
Nozzles:	Spring applications supported: CFF, VFFF, Twin Fluid and Air Inclusion

Best Use Advice - Spring

- Has a wide window of application up to crop GS32 inclusive
- Suitable for early spring control of wild oats (GS11-32 (optimal up to GS29)) and ryegrass (GS11-29) where:
 - Control was missed in the autumn
 - Pre-emergent product was not applied or was ineffective
 - Grassweeds have germinated in the spring
- Can apply at 210 g/ha for control of wild oats up to GS29
- Suitable for control of spring germinating bromes (GS11-24)
 - Optimally after effective pre-emergence
- Apply Broadway Star at 265 g/ha with an approved adjuvant:
 - Abacus, Activator 90, Bio-syl, Buzz, Drill, Cogent, Headland Diamond, Headland Fortune, Intracrop Felix, Intracrop Warrior, Kantor, Kinetic, Roller, Spartan, Spryte Aqua, Tonto, Torpedo II, Viking
- Apply during periods of active growth
- For optimal grassweed control use as part of a programme and follow WRAG guidelines
- No cultivation restrictions prior to sowing following crops:
 - Autumn: Winter wheat, barley, oats, triticale and rye, oilseed rape, grass, winter beans, brassica plants as transplants
 - Spring: Spring wheat, barley, oats, triticale, rye and oilseed rape, sugar beet, potatoes, grass, clover (as part of a grass/clover ley), beans, peas, maize, linseed, brassica plants as transplants, carrots, parsnips
- Rainfast 1 hour

TANK MIXING

- Broadway Star + adjuvant is physically compatible with a range of fungicides including products containing chlorothalonil
- Can be used with chlormequat PGRs at half dose rate. Other PGRs or higher doses require a 7 day interval
- Refer to <http://uk.dowagro.com/tank-mix-information> for further information

Weed Spectrum

- The market standard for **ryegrass** control (GS11-29) and **brome** control (GS11-24) 265 g/ha
- Outstanding **wild oat** control (GS11-29) 210 g/ha

	210 g/ha	265 g/ha		210 g/ha	265 g/ha		210 g/ha	265 g/ha
Bur-chervil		8 etl	Forget-me-not		BFSE	Redshank		6 etl
Black bindweed		4 etl	Geranium spp.	8 etl		Pale persicaria		6 etl
Charlock	6 etl		Groundsel		BFBV	Shepherd's purse		6 etl
Chickweed	100 mm		Hemp-nettle		4 etl	Shepherd's needle		BFSE
Common field speedwell	6 etl		Ivy-leaved speedwell	6 etl		Volunteer linseed		6 etl
Cleavers	150 mm		Knotgrass		2 etl; MS 6 etl	Volunteer beans		BFBV
Common vetch		6 etl	Mayweeds	8 etl		Volunteer oilseed rape	6 etl	
Corn marigold		4 etl	Mouse-ear chickweed		MS 100 mm	Volunteer peas		6 etl
Creeping thistle (from seed)		MS 6 etl	Mugwort		MS 150 mm	Wild radish		6 etl
Field pansy	4 etl		Poppy	6 etl				

Key: **bold** - label weeds; BFBV - Before flower buds visible; BFSE - Before stem extension; etl - established true leaves; MS - Moderately susceptible
 Non-Label weeds listed as an indication of the effect that would be expected to be achieved based on limited data.

