

## SAFETY PRECAUTIONS

### Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:  
**WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD)** when handling the concentrate. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.  
**DO NOT BREATHE SPRAY.**

**WHEN USING DO NOT EAT, DRINK OR SMOKE.**  
**WASH CONCENTRATE** from skin and eyes immediately.  
**WASH HANDS AND EXPOSED SKIN** before eating and drinking and after work.

### Environmental protection:

**DO NOT CONTAMINATE WATER** with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

### Storage and disposal:

**KEEP OUT OF REACH OF CHILDREN.**  
**KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.**  
**KEEP IN ORIGINAL CONTAINER**, tightly closed in a safe place.  
**RINSE CONTAINER THOROUGHLY** by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.  
**DO NOT RE-USE CONTAINER** for any purpose.



This label is compliant with the CPA Voluntary Initiative Guidance

 **Dow AgroSciences**



# GF-184

## HERBICIDE

Product Registration Number: MAPP 10878

A suspension emulsion formulation containing

100 g/litre fluroxypyr methyl heptyl ester and 2.5 g/litre florasulam.

A post-emergence herbicide for use on all varieties of **WINTER AND SPRING WHEAT, BARLEY AND OATS** for the control of a range of **BROAD-LEAVED WEEDS**.


The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

**READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.**

**PROTECT FROM FROST.**  
**SHAKE WELL BEFORE USE.**

## 5 Litres e

**Dow AgroSciences Limited**  
CPC2 Capital Park, Fulbourn,  
Cambridge, CB21 5XE.  
Telephone: (01462) 457272  
Fax: (01462) 426605  
24 Hour Emergency  
Telephone Number: +44 (0) 1553 761 251



**Product Identifier according to Art.18 of Reg. (EC) No 1272/2008 (CLP):** GF-184; Hydrocarbons, C9, aromatics

**WARNING**  
**CAUSES SKIN IRRITATION**  
**MAY CAUSE AN ALLERGIC SKIN REACTION**  
**CAUSES SERIOUS EYE IRRITATION**  
**MAY CAUSE RESPIRATORY IRRITATION**  
**MAY CAUSE DROWSINESS OR DIZZINESS**  
**VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS**

Wear protective gloves/clothing/eye/face protection.  
**IF ON SKIN:** Wash with plenty of soap and water.  
**IF INHALED:** Remove to fresh air and keep at rest in a position comfortable for breathing.  
**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.  
Do NOT induce vomiting.  
Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.  
**To avoid risks to human health and the environment, comply with the instructions for use.**

### IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops	Maximum Individual Dose	Maximum Total Dose	Latest Time of Application
Winter wheat, winter barley	1.8 litres product per hectare	1.8 litres product per hectare per crop	Before flag leaf sheath opening stage (up to Zadoks 45 inclusive)
Winter oats	1.8 litres product per hectare	1.8 litres product per hectare per crop	Before second node detectable stage (up to Zadoks 31 inclusive)
Spring wheat, spring barley	1.5 litres product per hectare	1.5 litres product per hectare per crop	Before flag leaf sheath extended stage (up to Zadoks 39 inclusive)
Spring oats	1.5 litres product per hectare	1.5 litres product per hectare per crop	Before second node detectable stage (up to Zadoks 31 inclusive)


### Other Specific Restrictions:

For autumn planted crops a maximum total dose of 3.75 g of florasulam must be observed for applications made between crop emergence in the year of planting and February 1<sup>st</sup> in the year of harvest.

The total amount of florasulam applied to a cereal crop must not exceed 7.5 g.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

P001987231611



## DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

## GENERAL INFORMATION

GF-184 is mainly absorbed through the foliage of weeds. The ideal timing for application is when the weeds are small and actively growing.

## NOTES

Do not spray when crops are under stress from cold, drought, pest damage, nutrient deficiency etc.

Do not roll or harrow 7 days before or after application.

Take extreme care to avoid drift onto crops and non-target plants outside the target area.

## WEED SUSCEPTIBILITY AND MAXIMUM SIZE CONTROLLED

Weed	Size Controlled
Cleavers*	Flowering
Chickweed	Flowering
Mayweed species	Flowering

\*Cleavers which germinate after application will not be controlled

## CROPS

GF-184 can be used on all varieties of winter and spring wheat, barley and oats.

## SOIL

GF-184 may be used on all soil types.

## TIMING

### Winter wheat, winter barley

GF-184 should be applied from 3 leaves (Zadoks 13) up to before flag leaf sheath opening stage (Zadoks 45).

### Spring wheat, spring barley

GF-184 should be applied from 3 leaves (Zadoks 13) up to and including before flag leaf sheath extending stage (Zadoks 39).

### Winter oats, spring oats

GF-184 should be applied from 3 leaves (Zadoks 13) up to and including the before second node detectable stage (Zadoks 31).

## RATES OF USE

One application of up to 1.8 litres/ha in winter crops or 1.5 litres/ha in spring crops will control all susceptible emerged weeds. A split application may be applied up to a maximum total dose of 1.8 litres/ha (winter crops) or 1.5 litres/ha (spring crops) where weed germination takes place over an extended period.

## APPLICATION

GF-184 may be applied through tractor-mounted hydraulic sprayers providing they are in good working order and have been calibrated according to the manufacturers' recommendations.

Do not apply through CDA applicators.

## WATER VOLUME

Apply GF-184 in 150 to 250 litres of water per hectare. The lower volume must only be used in open crops on small weeds.

## FOLLOWING CROPS

Crops that can be sown in the year of treatment with GF-184:

Cereals, oilseed rape, field beans, vegetable brassicas as transplants, grass.

\*Vigour reductions may be seen in following crops of oilseed rape after a dry summer.

This will be outgrown and will not result in yield loss.

Crops that can be sown the calendar year following treatment with GF-184:

Cereals, oilseed rape, field beans, carrots, vegetable brassicas as transplants, grass, linseed, peas, sugar beet, potatoes, maize, clover (for use in grass/clover mixtures).

## JOINT APPLICATION

A joint application is the use of a product in tank mixture or sequence with another product.

**IMPORTANT NOTE:** Joint applications should only be made within the label recommendations of every product in the application.

Only one other product with an ALS inhibitor mode of action may be applied to a cereal crop treated with GF-184. However a further application of GF-184 or another product containing florasulam may also be made **providing the maximum total dose of florasulam is not exceeded**¹.

GF-184 may be applied in joint application to the same cereal crop with one of the following ALS products:

Absolute	Galaxy¹
Alias	GF-184¹
Alias SX	GF-2070
Ally Express	Gropper SX
Ally Max SX	Harmony M SX
Ally SX	Harmony SX
Answer SX	Hatra
Avocet	Hiker¹
Avro SX	Horus
Biplay SX	Hunter¹
Broadway Star¹	Inka SX
Broadway Sunrise	iodosulfuron-methyl sodium (MAPP 12364, 15174, 16173)
Bullion	Jubilee SX
Calibre SX	Lancer
Chimera SX	Lexus Class
Concert SX	Lexus Millennium
Cleancrop Mondial	Lexus SX
Dakota¹	Lorate
Ductis SX	mesosulfuron-methyl + iodosulfuron-methyl sodium (MAPP 12049, 12478)
Excalibur	mesosulfuron-methyl (MAPP 11218)
Exceed SX	mesosulfuron-methyl + iodosulfuron-methyl sodium + diflufenican (MAPP 12695, 16149)
Finish SX	Mozaic SX

Octavian	Slalom¹
Oklar SX	Spitfire¹
Oriel 50SX	Staka SX
Parana	Starane Gold¹
Pennant	Starane Vantage¹
Pinnacle	Starane XL¹
Presite SX	Taxi
Quantum SX	Thor
Ratio SX	Traton SX
Refine Max SX	Triad
Seduce	Unite
Simba SX	

¹ The maximum total dose of florasulam applied to the crop must not exceed 7.5 g. For autumn planted crops a maximum total dose of 3.75 g of florasulam, must be observed for applications made between crop emergence in the year of planting and February 1<sup>st</sup> in the year of harvest.

Apart from these specific joint applications GF-184 must NOT be applied with any other product containing an ALS-inhibitor, for example amidosulfuron or triasulfuron.

## CROP FAILURE

In the event of a crop failure in the spring after treatment with GF-184, only the following crops may be planted: spring wheat, spring barley, spring oats, maize or ryegrass.

## MIXING

Half fill the spray tank with water and add the required amount of GF-184. Fill up the spray tank, agitating continuously to ensure thorough mixing, and maintain agitation until spraying is complete. Use only clean water for mixing.

## SPRAY QUALITY

Apply GF-184 as a MEDIUM spray as defined by the BCPC system.

## TANK MIXTURES

Where tank mixes are used, and unless directed otherwise, the preferred order of addition of products to the spray tank is as follows: water dispersible granules, wettable powders, suspension concentrates (flowables), emulsifiable concentrates, solution concentrates. Each product should be added to the half-full sprayer and be fully dispersed before the addition of the next product.

## TANK CLEANING

To avoid subsequent injury to crops other than cereals, all spraying equipment must be thoroughly cleaned both inside and out, using All Clear Extra spray cleaner as follows:

1. Immediately after spraying, drain tank completely. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
3. Half fill tank with clean water and add All Clear Extra at the recommended rate. Agitate and then briefly flush the booms and hoses with the cleaning solution. Top up with water making sure the tank is completely full and allow to stand for 15 minutes with agitation. Flush the booms and hoses and drain tank completely.
4. Nozzles and filters should be removed and cleaned separately with All Clear Extra solution containing 50 ml of All Clear Extra per 10 litres of water.

5. Rinse the tank with clean water and flush through the booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
6. For disposal of washings, follow Code of Practice for Using Plant Protection Products. Do not spray onto sensitive crop or land intended for cropping with sensitive crop.

**Note:** If it is not possible to drain the tank completely, step 3 must be repeated before going onto step 4.

#### **TRADEMARK ACKNOWLEDGEMENTS**


Broadway, Galaxy, Hiker, Hunter, Slalom, Spitfire, Starane, Sunrise and Vantage are trademarks of Dow AgroSciences LLC.

All other brand names are trademarks of other manufacturers for which proprietary rights may exist.

#### **DOW AGROSCIENCES CONDITIONS OF SUPPLY**

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

**To access the Safety Data Sheet for this product scan the QR code or use the weblink below:**



<http://uk.dowagro.com/wp-content/uploads/GF-184-SDS.pdf>

**Alternatively contact your supplier.**

SPECIMEN

**THIS PAGE HAS BEEN LEFT  
BLANK INTENTIONALLY**

**SPECIMEN**

## SAFETY PRECAUTIONS

### Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:  
WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.  
DO NOT BREATHE SPRAY.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin and eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

### Environmental protection:

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

### Storage and disposal:

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely. DO NOT RE-USE CONTAINER for any purpose.



Dow AgroSciences



# GF-184

## HERBICIDE

Product Registration Number: MAPP 10878

A suspension emulsion formulation containing 100 g/litre fluroxypyr methyl heptyl ester and 2.5 g/litre florasulam.

A post-emergence herbicide for use on all varieties of WINTER AND SPRING WHEAT, BARLEY AND OATS for the control of a range of BROAD-LEAVED WEEDS.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.


**READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.**

**PROTECT FROM FROST.  
SHAKE WELL BEFORE USE.**

## 5 Litres e

Dow AgroSciences Limited  
CPC2 Capital Park, Fulbourn,  
Cambridge, CB21 5XE.  
Telephone: (01462) 457272  
Fax: (01462) 426605  
24 Hour Emergency  
Telephone Number: +44 (0) 1553 761 251

This label is compliant with the  
CPA Voluntary Initiative Guidance



Product Identifier according to Art.18 of Reg. (EC) No 1272/2008

[CLP]: GF-184; Hydrocarbons, C9, aromatics

### WARNING

**CAUSES SKIN IRRITATION**

**MAY CAUSE AN ALLERGIC SKIN REACTION**

**CAUSES SERIOUS EYE IRRITATION**

**MAY CAUSE RESPIRATORY IRRITATION**

**MAY CAUSE DROWSINESS OR DIZZINESS**

**VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS**

Wear protective gloves/clothing/eye/face protection.

IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Do NOT induce vomiting.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

### IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops	Maximum Individual Dose	Maximum Total Dose	Latest Time of Application
Winter wheat, winter barley	1.8 litres product per hectare	1.8 litres product per hectare per crop	Before flag leaf sheath opening stage (up to Zadoks 45 inclusive)
Winter oats	1.8 litres product per hectare	1.8 litres product per hectare per crop	Before second node detectable stage (up to Zadoks 31 inclusive)
Spring wheat, spring barley	1.5 litres product per hectare	1.5 litres product per hectare per crop	Before flag leaf sheath extended stage (up to Zadoks 39 inclusive)
Spring oats	1.5 litres product per hectare	1.5 litres product per hectare per crop	Before second node detectable stage (up to Zadoks 31 inclusive)

### Other Specific Restrictions:

For autumn planted crops a maximum total dose of 3.75 g of florasulam must be observed for applications made between crop emergence in the year of planting and February 1<sup>st</sup> in the year of harvest.

The total amount of florasulam applied to a cereal crop must not exceed 7.5 g.

**READ THE LABEL BEFORE USE. APPLY THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

