

SAFETY PRECAUTIONS

Operator Protection:

WHEN USING DO NOT EAT, DRINK OR SMOKE.

Storage and Disposal:

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.
KEEP OUT OF REACH OF CHILDREN.
KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

Environmental protection:

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Product Identifier according to Art.18 of Reg. (EC) No 1272/2008 [CLP]:

Galera®



Toxic to aquatic life with long lasting effects

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for empty clean triple rinsed containers which can be disposed of as non-hazardous waste.

To avoid risks to man and the environment, comply with the instructions for use

Product Registration Number: MAPP 16413/PCS No. 05188

IMPORTANT INFORMATION.

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crop:	Oilseed rape (winter)
Maximum Individual Dose per Crop:	0.35 litre product per hectare
Maximum Total Dose per Crop:	0.35 litre product per hectare
Latest Time of Application:	Before flower buds visible above the crop canopy

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

© Trademark of The Dow Chemical Company ("Dow") or an Affiliated Company of Dow



Dow AgroSciences



Galera®

HERBICIDE

Product Registration Number: MAPP 16413 / PCS no. 05188

A soluble concentrate formulation containing 67 g/L picloram (present as 84 g/L picloramolamine) and 267 g/L clopyralid (present as 352 g/L clopyralid olamine).

A post-emergence herbicide for use on all varieties of WINTER OILSEED RAPE for the control of CLEAVERS and MAYWEEDS.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work. (UK only)

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.

SHAKE WELL BEFORE USE

PROTECT FROM FROST

Pack size: 1 Litre e

Dow AgroSciences Limited

CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE.

Telephone: (01462) 457272 Fax: (01462) 426605

24 Hour Emergency Telephone Number: +44 (0) 1553 761 251

PROFESSIONAL USE ONLY

Triple Rinse Containers, Puncture and Invert to Dry at time of Use



This label is compliant with the CPA Voluntary Initiative Guidance (UK Only).

P 002389401705



DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

GENERAL INFORMATION

GALERA® is a post-emergence herbicide for use on all varieties of winter oilseed rape for the control of cleavers and mayweeds. GALERA is mainly absorbed through the foliage of weeds and translocated to other plant parts. The ideal timing for application is when the weeds are small and actively growing.

FOLLOWING CROPS

Following application of Galera the following restrictions on planting the following crops must be observed:

Wheat, barley, oats, maize, oilseed rape: 4 months (120 days)

All other crops: 12 months (1 year)

Ploughing or thorough cultivation should be undertaken prior to planting leguminous crops (e.g. field beans and peas).

CROP FAILURE

In the event of crop failure of a spring treated GALERA crop, only the following crops may be planted: spring oilseed rape, spring wheat, spring barley, spring oats, maize or ryegrass (provided ploughing or thorough cultivation is undertaken prior to replanting).

NOTES

Do not use any plant material treated with GALERA for composting or mulching.

Do not use manure from animals fed on crops treated with GALERA for composting.

GALERA residues in plant tissues which have not completely decayed may affect succeeding susceptible crops.

If treated crop remains have not fully decayed by the time of planting following crops then avoid planting: peas, beans and other legumes; carrots and Umbellifers; potatoes; lettuce and other Compositae; glasshouse and protected crops.

Chop and incorporate all treated plant remains in the early autumn (or as soon as possible after harvest) to release any residues into the soil, where they are more quickly broken down. Following good agronomic practice ensure that stubble and straw (including farmyard manure and digestate) and other treated plant remains have completely decayed before planting susceptible crops.

Do not spray when crops are under stress from cold, drought, pest damage, nutrient deficiency etc.

Do not roll or harrow 7 days before or after application.

Take extreme care to avoid drift onto crops and non-target plants outside the target area.

Contract agents should be consulted before using on crops grown for seed.

WEED SUSCEPTIBILITY

To achieve optimum levels of weed control, good spray coverage of the target weed is required.

Weed Species	Weed Susceptibility	Maximum Growth Stage
Scentless mayweed	Susceptible	150 mm across/high
Scented mayweed	Susceptible	150 mm across/high
Cleavers*	Moderately susceptible	150 mm high/across

* Cleavers which germinate after application will not be controlled

CROPS

GALERA can be used on all varieties of winter oilseed rape.

TIMING

GALERA can be applied in the spring from the beginning of March (BBCH 20) up to but before flower buds are visible above the crop canopy (BBCH50).

RATES OF USE

One application of 0.35 litre/ha will control all susceptible emerged weeds.

WATER VOLUME

Apply GALERA in 200 litres of water per hectare.

SOIL

GALERA may be used on all soil types.

APPLICATION

GALERA may be applied through tractor-mounted hydraulic sprayers providing they are in good working order and have been calibrated according to the manufacturer's recommendations.

Do not apply through CDA applicators.

MIXING

Half fill the spray tank with water and add the required amount of GALERA. Fill up the spray tank, agitating continuously to ensure thorough mixing, and maintain agitation until spraying is complete. Use only clean water for mixing.

SPRAY QUALITY

Apply GALERA as a MEDIUM spray as defined by the BCPC system.

TANK CLEANING

Thoroughly wash all spraying and measuring equipment with water immediately after use.

To avoid subsequent injury to crops, all spraying equipment must be thoroughly cleaned both inside and out after an application of GALERA.

1. Immediately after spraying, drain tank completely. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
3. Half fill tank with clean water. Agitate and then briefly flush the boom and hoses. Top up with water making sure the tank is completely full and allow to stand for 15 minutes with agitation. Flush the boom and hoses and drain tank completely.
4. Nozzles and filters should be cleaned separately and removed if necessary.
5. For disposal of washings, follow local regulations. Do not spray onto sensitive crop or land intended for cropping with sensitive crop.

Note: If it is not possible to drain the tank completely, step 3 must be repeated before going onto step 4.

DOW AGROSCIENCES CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

UK:

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



<http://uk.dowagro.com/wp-content/uploads/GALERA-SDS.pdf>

Alternatively contact your supplier

REPUBLIC OF IRELAND:

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



<http://uk.dowagro.com/wp-content/uploads/GALERA-IRL-SDS.pdf>

Alternatively contact your supplier

SPECIMEN

**THIS PAGE HAS BEEN LEFT
BLANK INTENTIONALLY**

SPECIMEN

SAFETY PRECAUTIONS

Operator Protection:

WHEN USING DO NOT EAT, DRINK OR SMOKE.

Storage and Disposal:

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.
KEEP OUT OF REACH OF CHILDREN.
KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

Environmental protection:

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Product Identifier according to Art.18 of Reg. (EC) No 1272/2008 [CLP]:
Galera®



Toxic to aquatic life with long lasting effects

Dispose of contents / container to a licensed hazardous-waste disposal contractor or collection site except for empty clean triple rinsed containers which can be disposed of as non-hazardous waste.

To avoid risks to man and the environment, comply with the instructions for use
Product Registration Number: MAPP 16413/PCS No. 05188

IMPORTANT INFORMATION.

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crop:	Oilseed rape (winter)
Maximum Individual Dose per Crop:	0.35 litre product per hectare
Maximum Total Dose per Crop:	0.35 litre product per hectare
Latest Time of Application:	Before flower buds visible above the crop canopy

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

© Trademark of The Dow Chemical Company ("Dow") or an Affiliated Company of Dow



Dow AgroSciences



Galera®

HERBICIDE

Product Registration Number: MAPP 16413 / PCS no. 05188

A soluble concentrate formulation containing 67 g/L picloram (present as 84 g/L picloramolamine) and 267 g/L clopyralid (present as 352 g/L clopyralid olamine).

A post-emergence herbicide for use on all varieties of WINTER OILSEED RAPE for the control of CLEAVERS and MAYWEEDS.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work. (UK only)

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.

SHAKE WELL BEFORE USE

PROTECT FROM FROST

Pack size: 1 Litre e

Dow AgroSciences Limited

CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE.

Telephone: (01462) 457272 Fax: (01462) 426605

24 Hour Emergency Telephone Number: +44 (0) 1553 761 251

PROFESSIONAL USE ONLY

Triple Rinse Containers, Puncture and Invert to Dry at time of Use



This label is compliant with the CPA Voluntary Initiative Guidance (UK Only).