

Pixxaro® EC

ARYLEX™ ACTIVE

TECHNICAL INFORMATION SHEET

Pixxaro® EC is a new herbicide for the control of broad-leaved weeds in winter cereals (wheat, durum wheat, spelt, barley, rye and triticale, and spring cereals (wheat, durum wheat, barley and rye).

Pixxaro EC is a **non-ALS**, spring-applied, wide spectrum broad-leaved weed herbicide, for cereals, containing ARYLEX™ Active. Regardless of weather conditions, Pixxaro EC provides robust control of highly competitive weeds such as cleavers chickweed, and poppy.

Why use Pixxaro EC?

- Provides robust weed control irrespective of the weather or weed growth pattern – if you can travel it will work
- Wide range of following crop options
- No cultivations required before sowing most crops
- Robust weed control: **chickweed, cleavers, fumitory, fat hen, dead nettle, poppy** and **cranesbill**
- Excellent multiway compatibility including: ACCase graminicides, ALS graminicides, PGRs Fungicides and Trace elements

Key facts

- Product Registration Number: MAPP 17545
- Active Ingredient: 12 g/litre Arylex + 280 g/litre fluroxypyr
- Pack size: 5.0 Litres
- Formulation: Emulsifiable Concentrate (Liquid)
- Maximum Individual Dose: 0.5 L/ha from 1st February (Winter cereals)
0.5 L/ha from 1st March (Spring cereals)
- Maximum Total Dose: 0.5 L/ha
- Application Timing: Winter cereals 1st February to 30th June GS13 to GS45
Spring cereals 1st March to 30th June GS13 to GS45
- Water Volumes: 100-400 L/ha
- Recommended Adjuvants: Non-ionic wetters and methylated seed oils
- Spray Quality: Medium as defined by BCPC
- Nozzles: Flat Fan, Variable Pressure Flat Fan, Pre-Orifice, Air Inclusion, Airtec
- Buffer Zone: 5m reducible buffer zone (1m dry ditches)



Dow AgroSciences

Solutions for the Growing World


Use plant protection products safely. Always read the label and product information before use. For further information including warning phrases and symbols refer to label. Dow AgroSciences Limited, CPC2 Capital Park, Fulbourn, Cambridge, CB21 5XE. Tel: +44 (0) 1462 457272.™ Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow. All other brand names are trademarks of other manufacturers for which proprietary rights may exist. PIXXARO® EC contains halauxifen-methyl (Arylex™ Active) and fluroxypyr. Technical Hotline: 0800 689 8899 | UKHotline@dow.com | uk.dowagro.com

Revised April 2017

Where Does Pixxaro EC fit?

Winter Cereals

- Tank mix with spring applied ALS graminicide
- Tank mix with T0 or T1 fungicide programme
- Compatible in multiway mixes with fungicides/PGRs/trace elements/herbicides
- For late season control of **Cleavers** (GS34-45 of the crop), use Pixxaro EC at 0.25L/ha



Spring Cereals

- Tank mix with ½ rate SU or dicamba product
- Tank mix with ACCase graminicide
- Apply mid to late tillering of the cereal crop
- Flexibility of tank mix partners including graminicides, PGRs & fungicides
- For late season control of **Cleavers** (GS34-45 of the crop), use Pixxaro EC at 0.25L/ha



More Information:

For product label, tank mix compatibility and more information on Pixxaro EC please see:

uk.dowagro.com/products/pixxaro-ec/

Weed Spectrum

| Weed | 0.375 L/ha | 0.5 L/ha |
|-------------------------|--------------|--------------------|
| Black Bindweed* | S<10 cm | S<15 cm |
| Black Nightshade | - | S<4 lf |
| Burr Chervil | T | T |
| Charlock | T | T |
| Chickweed* | S< 25 cm | S<flowering |
| Cleavers | b4 flowering | b4 flowering |
| Clover | - | S<4 lf |
| Cranesbill* | S<5 cm | S<10 cm |
| Docks, Seedling | - | S<4 lf, up to 5 cm |
| Fat Hen | S<15 cm | S<25 cm |
| Forget-me-not | MS<10 cm | S<5 cm |
| Fool's Parsley | - | S<4 lf |
| Fumitory* | S<20 cm | b4 flowering |
| Groundsel | T | MS |
| Hemp-nettle | S< 10 cm | b4 flowering |
| Henbit Deadnettle | S<flowering | S<flowering |
| Knotgrass* | MS | MS |
| Marigold | MS | MS |
| Mayweed | T | T |
| Nettle, Small | S | S |
| Orache | MS | MS |
| Pale Persicaria* | MS | MS |
| Pennycress | S<10 cm | S<10 cm |
| Pansy | T | T |
| Poppy* | S<10 cm | Rosette |
| Red Dead Nettle | S<18 cm | b4 flowering |
| Redshank* | S<3 lf | S<3 lf |
| Scarlet Pimpernel | MS | S<flowering |
| Shepherds Purse | S<10 cm | b4 flowering |
| Shepherds Needle | T | T |
| Speedwell, Common Field | T | S<4 lf, MS>4 lf |
| Speedwell, Ivy Leaved | MS<2 lf | S<4 lf, MS>4 lf |
| Spurrey | - | S<2 lf |
| Weed beet# | T | T |
| Wild carrot | MS<4 lf | S<4 lf |
| Wild Radish | T | T |
| Vol Beans | S<8 lf | b4 flowering |
| Vol Borage | | S<2 lf |
| Vol Linseed | | S<10 cm |
| Vol Oilseed Rape | T | T |
| Vol Peas | S<8 lf | b4 flowering |
| Vol Potatoes* | T | MS<20 cm |

Key: **bold** – label weeds, * – improved control with adjuvant, S – susceptible, MS – moderately susceptible, T – tolerant, # – <50% control

Non-label weeds listed as an indication of the effect that would be expected to be achieved based on limited data.