

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:
WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

DO NOT BREATHE SPRAY.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin and eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

Environmental protection:

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and disposal:

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely. DO NOT RE-USE CONTAINER for any purpose.



Product Registration Number: MAPP 13772

A suspension emulsion formulation containing 100 g/litre fluroxypyr methyl heptyl ester and 2.5 g/litre florasulam.

A post-emergence herbicide for use on all varieties of WINTER AND SPRING WHEAT, BARLEY AND OATS for the control of a range of BROAD-LEAVED WEEDS.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.

PROTECT FROM FROST.
SHAKE WELL BEFORE USE.

5 Litres e



3 362130 059868 >

Dow AgroSciences Limited
CPC2 Capital Park, Fulbourn, Cambridge,
CB21 5XE.
Telephone: (01462) 457272
Fax: (01462) 426605
24 Hour Emergency
Telephone Number: +44 (0) 1553 761 251



This label is compliant with the
CPA Voluntary Initiative Guidance

Product Identifier according to Art.18 of Reg. (EC) No 1272/2008 [CLP]:
Slalom®: Hydrocarbons, C9, aromatics

WARNING
CAUSES SKIN IRRITATION
MAY CAUSE AN ALLERGIC SKIN REACTION
CAUSES SERIOUS EYE IRRITATION
MAY CAUSE RESPIRATORY IRRITATION
MAY CAUSE DROWSINESS OR DIZZINESS
VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS

Wear protective gloves/clothing/eye/face protection.
IF ON SKIN: Wash with plenty of soap and water.
IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
Do NOT induce vomiting.
Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.
To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops	Maximum Individual Dose	Maximum Total Dose	Latest Time of Application
Winter wheat, winter barley	1.8 litres product per hectare	1.8 litres product per hectare per crop	Before flag leaf sheath opening stage (up to Zadoks 45 inclusive)
Winter oats	1.8 litres product per hectare	1.8 litres product per hectare per crop	Before second node detectable stage (up to Zadoks 31 inclusive)
Spring wheat, spring barley	1.5 litres product per hectare	1.5 litres product per hectare per crop	Before flag leaf sheath extended stage (up to Zadoks 39 inclusive)
Spring oats	1.5 litres product per hectare	1.5 litres product per hectare per crop	Before second node detectable stage (up to Zadoks 31 inclusive)

Other Specific Restrictions:

For autumn planted crops a maximum total dose of 3.75 g of florasulam must be observed for applications made between crop emergence in the year of planting and February 1st in the year of harvest.

The total amount of florasulam applied to a cereal crop must not exceed 7.5g.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

®Trademark of the Dow Chemical Company ('Dow') or an affiliated company of Dow



P 0 0 3 1 7 7 5 3 1 6 1 0



DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

GENERAL INFORMATION

SLALOM® is mainly absorbed through the foliage of weeds. The ideal timing for application is when the weeds are small and actively growing.

NOTES

Do not spray when crops are under stress from cold, drought, pest damage, nutrient deficiency etc.

Do not roll or harrow 7 days before or after application.

Take extreme care to avoid drift onto crops and non-target plants outside the target area.

WEED SUSCEPTIBILITY AND MAXIMUM SIZE CONTROLLED

Weed	Size Controlled
Cleavers*	Flowering
Chickweed	Flowering
Mayweed species	Flowering

*Cleavers which germinate after application will not be controlled

CROPS

SLALOM can be used on all varieties of winter and spring wheat, barley and oats.

SOIL

SLALOM may be used on all soil types.

TIMING

Winter wheat, winter barley

SLALOM should be applied from 3 leaves (Zadoks 13) up to before flag leaf sheath opening stage (Zadoks 45).

Spring wheat, spring barley

SLALOM should be applied from 3 leaves (Zadoks 13) up to and including before flag leaf sheath extending stage (Zadoks 39).

Winter oats, spring oats

SLALOM should be applied from 3 leaves (Zadoks 13) up to and including the before second node detectable stage (Zadoks 31).

RATES OF USE

One application of up to 1.8 litres/ha in winter crops or 1.5 litres/ha in spring crops will control all susceptible emerged weeds. A split application may be applied up to a maximum total dose of 1.8 litres/ha (winter crops) or 1.5 litres/ha (spring crops) where weed germination takes place over an extended period.

APPLICATION

SLALOM may be applied through tractor-mounted hydraulic sprayers providing they are in good working order and have been calibrated according to the manufacturers' recommendations. Do not apply through CDA applicators.

WATER VOLUME

Apply SLALOM in 150 to 250 litres of water per hectare. The lower volume must only be used in open crops on small weeds.

FOLLOWING CROPS

Crops that can be sown in the year of treatment with SLALOM:

Cereals, oilseed rape¹, field beans, vegetable brassicas as transplants, grass.

¹Vigour reductions may be seen in following crops of oilseed rape after a dry summer.

This will be outgrown and will not result in yield loss.

Crops that can be sown the calendar year following treatment with SLALOM:

Cereals, oilseed rape, field beans, carrots, vegetable brassicas as transplants, grass, linseed, peas, sugar beet, potatoes, maize, clover (for use in grass/clover mixtures).

JOINT APPLICATION

A joint application is the use of a product in tank mixture or sequence with another product.

IMPORTANT NOTE: Joint applications should only be made within the label recommendations of every product in the application.

Only one other product with an ALS-inhibitor mode of action may be applied to a cereal crop treated with SLALOM.

However a further application of SLALOM or another product containing florasulam may also be made providing the maximum total dose of florasulam is not exceeded¹.

SLALOM may be applied in joint application to the same cereal crop with one of the following ALS products:

Absolute	Exceed SX
Alias	Finish SX
Alias SX	Galaxy ¹
Ally Express	GF-184 ¹
Ally Max SX	GF-2070
Ally SX	Gropper SX
Answer SX	Harmony M SX
Avocet	Harmony SX
Avro SX	Hatra
Biplay SX	Hiker ¹
Broadway Star ¹	Horus
Broadway Sunrise	Inka SX
Bullion	iodosulfuron-methyl sodium (MAPP 12364, 15174, 16173)
Calibre SX	Jubilee SX
Chimera SX	Lancer
Concert SX	Lexus Class
Cleancrop Mondial	Lexus Millenium
Dakota ¹	Lexus SX
Ductis SX	Lorate
Excalibur	mesosulfuron-methyl + iodosulfuron-methyl sodium (MAPP 12049, 12478)

mesosulfuron-methyl (MAPP 11218)	Refine Max SX
mesosulfuron-methyl + iodosulfuron-methyl sodium + diflufenican (MAPP 12695, 16149)	Seduce
Mozaic SX	Simba SX
Octavian	Slalom ¹
Oklar SX	Spitfire ¹
Oriel 50SX	Staka SX
Parana	Starane Gold ¹
Pennant	Starane Vantage ¹
Pinnacle	Starane XL ¹
Presite SX	Taxi
Quantum SX	Thor
Ratio SX	Traton SX

¹The maximum total dose of florasulam applied to the crop must not exceed 7.5g. For autumn planted crops a maximum total dose of 3.75g of florasulam must be observed for applications made between crop emergence in the year of planting and February 1st in the year of harvest.

Apart from these specific joint applications SLALOM must NOT be applied with any other product containing an ALS-inhibitor, for example amidosulfuron or triasulfuron.

CROP FAILURE

In the event of a crop failure in the spring after treatment with SLALOM, only the following crops may be planted: spring wheat, spring barley, spring oats, maize or ryegrass.

MIXING

Half fill the spray tank with water and add the required amount of SLALOM. Fill up the spray tank, agitating continuously to ensure thorough mixing, and maintain agitation until spraying is complete. Use only clean water for mixing.

SPRAY QUALITY

Apply SLALOM as a MEDIUM spray as defined by the BCPC system.

TANK MIXTURES

Where tank mixes are used, and unless directed otherwise, the preferred order of addition of products to the spray tank is as follows: water dispersible granules, wettable powders, suspension concentrates (flowables), emulsifiable concentrates, solution concentrates. Each product should be added to the half-full sprayer and be fully dispersed before the addition of the next product.

TANK CLEANING

To avoid subsequent injury to crops other than cereals, all spraying equipment must be thoroughly cleaned both inside and out, using All Clear Extra spray cleaner as follows:

1. Immediately after spraying, drain tank completely. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
3. Half fill tank with clean water and add All Clear Extra at the recommended rate. Agitate and then briefly flush the booms and hoses with the cleaning solution. Top up with water making sure the tank is completely full and allow to stand for 15 minutes with agitation. Flush the booms and hoses and drain tank completely.

- Nozzles and filters should be removed and cleaned separately with All Clear Extra solution containing 50 ml of All Clear Extra per 10 litres of water.
- Rinse the tank with clean water and flush through the booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
- For disposal of washings, follow Code of Practice for the Using Plant Protection Products. Do not spray onto sensitive crop or land intended for cropping with sensitive crop.

Note: If it is not possible to drain the tank completely, step 3 must be repeated before going onto step 4.

TRADEMARK ACKNOWLEDGEMENTS

SLALOM, Broadway, Galaxy, Hiker, Hunter, Spitfire, Starane, Sunrise and Vantage are trademarks of Dow AgroSciences LLC.

All other brand names are trademarks of other manufacturers for which proprietary rights may exist.

DOW AGROSCIENCES CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

®Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



<http://uk.dowagro.com/wp-content/uploads/SLALOM-SDS.pdf>

Alternatively contact your supplier.

SPECIMEN

**THIS PAGE HAS BEEN LEFT
BLANK INTENTIONALLY**

SPECIMEN

SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:
WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.
DO NOT BREATHE SPRAY.
WHEN USING DO NOT EAT, DRINK OR SMOKE.
WASH CONCENTRATE from skin and eyes immediately.
WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

Environmental protection:

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and disposal:

KEEP OUT OF REACH OF CHILDREN.
KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.
KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.
RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.
DO NOT RE-USE CONTAINER for any purpose.

 **Dow AgroSciences**

Slalom[®]

HERBICIDE

Product Registration Number: MAPP 13772

A suspension emulsion formulation containing 100 g/litre fluroxypyr methyl heptyl ester and 2.5 g/litre florasulam.

A post-emergence herbicide for use on all varieties of WINTER AND SPRING WHEAT, BARLEY AND OATS for the control of a range of BROAD-LEAVED WEEDS.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.

**PROTECT FROM FROST.
SHAKE WELL BEFORE USE.**

5 Litres e

Dow AgroSciences Limited
CPC2 Capital Park, Fulbourn, Cambridge,
CB21 5XE.
Telephone: (01462) 457272
Fax: (01462) 426605
24 Hour Emergency
Telephone Number: +44 (0) 1553 761 251



This label is compliant with the
CPA Voluntary Initiative Guidance



Product Identifier according to Art.18 of Reg. (EC) No 1272/2008 [CLP]:
Slalom[®]; Hydrocarbons, C9, aromatics

WARNING

**CAUSES SKIN IRRITATION
MAY CAUSE AN ALLERGIC SKIN REACTION
CAUSES SERIOUS EYE IRRITATION
MAY CAUSE RESPIRATORY IRRITATION
MAY CAUSE DROWSINESS OR DIZZINESS
VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS**

Wear protective gloves/clothing/eye/face protection.
IF ON SKIN: Wash with plenty of soap and water.
IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
Do NOT induce vomiting.
Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.
To avoid risks to human health and the environment, comply with the instructions for use.

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops	Maximum Individual Dose	Maximum Total Dose	Latest Time of Application
Winter wheat, winter barley	1.8 litres product per hectare	1.8 litres product per hectare per crop	Before flag leaf sheath opening stage (up to Zadoks 45 inclusive)
Winter oats	1.8 litres product per hectare	1.8 litres product per hectare per crop	Before second node detectable stage (up to Zadoks 31 inclusive)
Spring wheat, spring barley	1.5 litres product per hectare	1.5 litres product per hectare per crop	Before flag leaf sheath extended stage (up to Zadoks 39 inclusive)
Spring oats	1.5 litres product per hectare	1.5 litres product per hectare per crop	Before second node detectable stage (up to Zadoks 31 inclusive)

Other Specific Restrictions:

For autumn planted crops a maximum total dose of 3.75 g of florasulam must be observed for applications made between crop emergence in the year of planting and February 1st in the year of harvest.

The total amount of florasulam applied to a cereal crop must not exceed 7.5g.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.



©Trademark of the Dow Chemical Company ("Dow") or an affiliated company of Dow