

**For Weed Control in Game Cover Crops**
**Key Facts**

Product Registration Number:	MAPP 16123
Active Ingredient:	200 g/litre clopyralid + 200 g/litre triclopyr
Pack Size:	3.0 Litres
Formulation:	Solution Concentrate
Maximum Individual Dose:	1.0 L/ha
Maximum Total Dose:	1.0 L/ha (one application per year)
Latest Timing:	After 1 <sup>st</sup> March but before end of October in year of application
Water Volumes:	200 – 400 L/ha
Spray Quality:	MEDIUM as defined by BCPC
LERAP:	None

When applying Thistlex to Environmental Stewardship Scheme options please check with your local scheme adviser

**Best Use Advice for Game Cover Crops**

Approval No.	Expiry Date	Crop
20142512	31/10/20	Outdoor crops grown for game cover.

- Use of Thistlex is permitted for use in game cover where no part of the plant is to be consumed by humans or used as food or feed for animals (excluding game birds).
- More information including use rates, timing etc. are given in the Extension of Authorisation Notice which is available from CRD website - <http://secure.pesticides.gov.uk/offlabels/search.asp>

**Game Cover species considered selective for use with Thistlex:**

- |                |                             |                                    |
|----------------|-----------------------------|------------------------------------|
| • Borage       | • Maize                     | • Sorghum                          |
| • Buckwheat    | • Millet (white, red, reed) | • Triticale and other cereals      |
| • Canary grass | • Mustard *                 | • Tyfon cabbage / stubble turnip * |
| • Kale         | • Phacelia *                | • Linseed *                        |

\* May be some crop damage resulting in loss of crop vigour of between 15% - 40% (mean of 3 trials).

NB Dow Shield 400 may be a safer alternative on some brassica type game covers – see Dow Shield 400 game cover technical information sheet for further information.

**Game Cover species controlled with Thistlex:**

- |                 |                    |             |
|-----------------|--------------------|-------------|
| • Chicory       | • Gold of pleasure | • Lupin     |
| • Fodder radish | • Quinoa           | • Sunflower |

**Following crops:** Do not plant winter beans in the same year as treatment with Thistlex. For sensitive spring crops (e.g. peas, beans, carrots, potatoes, tomatoes, lettuce, protected crops) don't spray Thistlex after 31<sup>st</sup> July.

**This extension of the authorised use provides for the use of Thistlex in respect of crops and situations, other than those included on the product label. No efficacy or phytotoxicity data have been assessed and as such the 'extension of use', is at all times done at the user's choosing, and the commercial risk is entirely theirs.**

**Weed Spectrum**

Annual Weeds	Control	Perennial Weeds	Control
Black bindweed	MS	Bramble	MT
Chickweed	MS	Clover	S
Corn chamomile	S	Dandelion	MT
Corn marigold	S	Docks	MT
Fat-hen	S	Nettles	MS
Groundsel	S	Ragwort	T
Mayweed	S	<b>Thistles</b> (creeping, spear)	S
		vetch	S

Key: **Bold** – label weed, S – susceptible, MS - moderately susceptible, MT – moderately tolerant, T - tolerant


**Dow AgroSciences**
*Solutions for the Growing World*